



Chitin (CBD-tag Affinity) Magnetic Beads

Catalog No. MBPP-06001, MBPP-06002, MBPP-06003

【Introduction】

Chitin Magnetic Beads are nano-superparamagnetic beads covalently coupled with chitin. With a fast magnetic response rate, high protein binding capacity and low non-specific binding, Chitin Magnetic Beads provide a rapid and efficient method to purify CBD(chitin-binding domain)-fusion proteins from cell culture supernatant. The beads are simply added to cell culture supernatant and CBD-fusion proteins will bind to the beads. After washing unbound proteins off, the CBD-fusion proteins can be eluted from the magnetic beads or the protein-bound magnetic beads can be directly used in downstream experiments (e.g. capturing target proteins, which bind to the immobilized CBD-fusion proteins, from crude cell lysates). The process can be completed manually or fully automated for high throughput applications.

【Product Specifications】

- **Diameter:** 500nm
- **pH stability:** pH 3-13
- **30min sedimentation rate:** <0.1%
- **Magnetic response rate:** >30emu/g
- **Solvent:** 20% ethanol
- **Binding capacity:** 60-100 μ g CBD-fusion proteins per mg magnetic beads

【Product Content】

<i>Catalog No.</i>	<i>Conc. (mg/ml)</i>	<i>Volume (ml)</i>	<i>Amount of Beads (mg)</i>
MBPP-06001	50	1	50
MBPP-06002	50	4	200
MBPP-06003	50	20	1000



Advanced BioChemicals (ABC) Advance Your Future

Phone: 1-678-827-9993 Fax: 1-509-696-4035 sales@advancedbiochemicals.com

【Purification Protocol】

The following protocol provides general guidelines for purification of CBP-fusion proteins using Chitin Magnetic Beads and may be modified by the user for specific applications. The protocol is scalable.

A. Additional material recommended

1. Binding/Washing Buffer: 20mM Tris-HCl, 0.5M NaCl, 1mM EDTA, 0.1% Tween-20, pH 8.0
2. A magnetic stand or a 96-well magnetic bead automation processor

B. Isolation of CBP-fusion proteins

1. Gently mix the magnetic beads thoroughly before use by repeated inversion.
2. Place 20 μ l of magnetic beads (1mg) into a 1.5ml sterile microcentrifuge tube.
3. Place the tube on a magnetic stand, collect the beads and discard the supernatant.
4. Wash the beads twice with Binding/Washing Buffer (500 μ l each time) by magnetic separation. Collect the beads and discard the supernatant.
5. Add 200-500 μ l of cell culture supernatant to the beads, mix thoroughly and incubate for 1.0hr at 4°C on a rotator.
6. Collect the beads with a magnet and save the supernatant for analysis if desired.
7. Wash the protein-coupled beads three times with Binding/Washing Buffer (500 μ l each time) by magnetic separation.
8. The CBD-fusion proteins can be eluted from the magnetic beads (e.g. boiling in 30 μ l of SDS-PAGE reducing sample buffer for 5min).
9. Or the protein-bound magnetic beads can be directly used in downstream experiments (e.g. capturing target proteins, which bind to the immobilized CBD-fusion proteins, from crude cell lysates).

【Storage】

Stored at 2-8°C, 2 years.

【Manufacturer】

Avanbio, ABA-06001, ABA-06002, ABA-06003

Advanced BioChemicals, LLC
P.O. Box 465066
Lawrenceville, GA 30042
USA

www.advancedbiochemicals.com
sales@advancedbiochemical.com
Phone: 1-678-827-9993
Fax: 1-509-696-4035